

All correspondence to:

**The Breastfeeding Network**  
**PO Box 11126, Paisley PA2 8YB**  
Admin Tel: 0844 412 0995  
e-mail: [admin@breastfeedingnetwork.org.uk](mailto:admin@breastfeedingnetwork.org.uk)  
[www.breastfeedingnetwork.org.uk](http://www.breastfeedingnetwork.org.uk)

## Breastfeeding and podiatry. local anaesthetics and liquefied phenol for ingrowing toenails

*The information provided is taken from various reference sources. It is provided as a guideline. No responsibility can be taken by the author or the Breastfeeding Network for the way in which the information is used. Clinical decisions remain the responsibility of medical and breastfeeding practitioners. The data presented here is intended to provide some immediate information but cannot replace input from professionals.*

### Ingrowing toenail surgery involving local anaesthetic and phenol does not necessitate interruption of breastfeeding

The available research evidence of safety from studies, for the administration of phenol to ingrowing toenails during lactation is limited. However, from the study of pharmacokinetic data, the risk is low and the advantages of ongoing breastfeeding are greater. The risk from the use of local anaesthetics during lactation is negligible.

The passage of phenol into breastmilk has not been extensively studied. However, absorption of topical products into breastmilk is restricted (1) so the limited quantity of liquid phenol applied to the nailbed during the procedure is unlikely to penetrate into breastmilk in any significant quantities. The passage across the different biological membranes/systems which have to be traversed before absorption from breastmilk, suggest that the application is unlikely to affect the breastfed baby.

The EPA (US Environmental protection agency) (2) has determined that exposure to phenol in drinking water at a concentration of 6 milligrams per litre (mg/L) for up to 10 days is not expected to cause any adverse effects in a child.

Phenol is considered to be quite toxic to humans via oral exposure but studies represent chronic rather than single episode exposure (3). Solutions containing phenol should not be applied to **large** areas of skin or wounds since sufficient phenol may be absorbed to give rise to toxic symptoms. This is not applicable in the use in the treatment of ingrowing toe-nails.

There is no information on levels of phenol in human breast milk (4). No mention is made of use in lactation in the Cochrane reviews (5,6).

Local anaesthetics are poorly bio available and have a short half- life and should not preclude normal breastfeeding after the procedure. They are widely used in other areas of medicine during breastfeeding e.g dental procedures, suturing post-natally and in other surgical procedures. (5,6,7,8,9) .

**To speak to a Breastfeeding Supporter call the National Breastfeeding Helpline 0300 100 0212**

Calls to 0300 numbers cost no more than calls to UK numbers starting 01 and 02 and will be part of any inclusive minutes that apply to your provider and call package

The Breastfeeding Network is a Company Limited by Guarantee Registered in Scotland Company No. 330639  
Registered office Whitelaw Wells, 9 Ainslie Place, Edinburgh, EH3 6AT  
The Breastfeeding Network is a Registered Scottish Charity No SC027007

©Wendy Jones PhD, MRPharmS and the Breastfeeding Network May 2017

To find your nearest Breastfeeding Supporter call the **Supporterline 0300 100 0210**

This information was accepted by the College of Podiatry in March 2017 and forms part of their current guidelines.

### References

1. Stoukides C Topical Medications and Breastfeeding. J Hum Lact 1993; 9(3) :185-7)
2. EPA Public Health Statement 2016 [www.epa.gov/sites/production/files/2016-09/documents/phenol.pdf](http://www.epa.gov/sites/production/files/2016-09/documents/phenol.pdf)
3. EPA Hazard Study Phenol 2016 [www.epa.gov/sites/production/files/2016-09/documents/phenol.pdf](http://www.epa.gov/sites/production/files/2016-09/documents/phenol.pdf)
4. Toxicological Profile For Phenol [www.atsdr.cdc.gov/ToxProfiles/tp115-c1.pdf](http://www.atsdr.cdc.gov/ToxProfiles/tp115-c1.pdf)
5. Cochrane review. Treatment for ingrowing toenails 2012  
[http://www.cochrane.org/CD001541/SKIN\\_treatments-for-ingrowing-toenails](http://www.cochrane.org/CD001541/SKIN_treatments-for-ingrowing-toenails)
6. Cochrane clinical answers What are the benefits and harms of interventions for ingrowing toenails?  
<http://cochraneclinicalanswers.com/doi/10.1002/cca.484/full>
7. Hale T Medications and Mothers Milk Hale Publishing 2016
8. Jones W Breastfeeding and Medication Routledge 2013
9. Lactmed ToxNet database <https://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?LACTMED>



To speak to a Breastfeeding Supporter call the **National Breastfeeding Helpline 0300 100 0212**

*Calls to 0300 numbers cost no more than calls to UK numbers starting 01 and 02 and will be part of any inclusive minutes that apply to your provider and call package*

*The Breastfeeding Network is a Company Limited by Guarantee Registered in Scotland Company No. 330639  
Registered office Whitelaw Wells, 9 Ainslie Place, Edinburgh, EH3 6AT  
The Breastfeeding Network is a Registered Scottish Charity No SC027007*

©Wendy Jones PhD, MRPharmS and the Breastfeeding Network May 2017