

All correspondence to:

The Breastfeeding Network
PO Box 11126, Paisley PA2 8YB
Admin Tel: 0844 412 0995
e-mail: admin@breastfeedingnetwork.org.uk
www.breastfeedingnetwork.org.uk

Shingles and Breastfeeding

The information provided is taken from various reference sources. It is provided as a guideline. No responsibility can be taken by the author or the Breastfeeding Network for the way in which the information is used. Clinical decisions remain the responsibility of medical and breastfeeding practitioners. The data presented here is intended to provide some immediate information but cannot replace input from professionals.

Aciclovir can be taken by a breastfeeding mother with shingles together with paracetamol and a non-steroidal drug e.g. ibuprofen

Shingles, also known as herpes zoster, is a painful skin rash caused by the reactivation of the chickenpox virus (varicella-zoster virus) in people who have previously had chickenpox. Early symptoms can include a headache, burning, tingling, numbness or itchiness of the skin in the affected area (normally the area is in a line often across the tummy, back or face), a feeling of being generally unwell accompanied by a temperature.

Many mothers with shingles will be prescribed a course of antiviral tablets lasting seven days, usually acyclovir 800mg .

Commonly prescribed antiviral medicines include aciclovir, famciclovir and valaciclovir. These medications can't kill the shingles virus, but can help stop the virus multiplying reducing the severity of the symptoms and lowering the risk of post-herpetic neuralgia which can last several months. Medicines are most effective when taken within 72 hours of symptoms so it is important to consult a healthcare professional quickly. You may also need to take painkilling medication such as paracetamol and ibuprofen (see factsheet on analgesics and breastfeeding).

If the rash is only on your body and can be covered by clothing, there's little risk of passing the infection on to others even before the lesions have dried.

- **Aciclovir** (Zovirax ®) can be given during lactation as it achieves less than 1% of the paediatric licensed dose. Oral bio-availability of aciclovir is limited. It is widely used to treat neonates and its safety is well established Lau 1987, Frenkel 1991, Meter 1988, Taddio 1994)
- **Famciclovir** (Famvir ®) has greater bio-availability than aciclovir but data is not available on its transfer into breastmilk. It has no particular benefits over aciclovir and is more expensive

To speak to a Breastfeeding Supporter call the National Breastfeeding Helpline 0300 100 0212

Calls to 0300 numbers cost no more than calls to UK numbers starting 01 and 02 and will be part of any inclusive minutes that apply to your provider and call package

The Breastfeeding Network is a Company Limited by Guarantee Registered in Scotland Company No. 330639
Registered office Whitelaw Wells, 9 Ainslie Place, Edinburgh, EH3 6AT
The Breastfeeding Network is a Registered Scottish Charity No SC027007

©Wendy Jones PhD, MRPharmS and the Breastfeeding Network January 2017

so unless there are compelling reasons why this drug should be used, aciclovir would be preferred during lactation. It is not used in children.


- **Valacyclovir** (Valtrex®) is a prodrug that is rapidly converted to acyclovir in the body.

References

- Lau RJ, Emery MG, Galinsky RE, Unexpected accumulation of acyclovir in breastmilk with estimation of infant exposure, *Obstet Gynecol*, 1987;69:468–71.
- Frenkel LM, Brown ZA, Bryson YJ, Corey L, Unadkat JD, Hensleigh PA, Arvin AM, Prober CG, Connor JD, Pharmacokinetics of acyclovir in the term human pregnancy and neonate, *Am J Obstet Gynecol*, 1991;164:569–76.
- Meyer LJ, de Miranda P, Sheth N, Spruance S, Acyclovir in human breastmilk, *Am J Obstet Gynecol*, 1988;158:586–8.
- Taddio A, Klein J, Koren G, Acyclovir excretion in human breastmilk, *Ann Pharmacother*, 1994;28:585–7
-

Bibliography

- Jones W *Breastfeeding and Medication* Routledge 2013
- Hale T *Medications and Mothers Milk* – online access January 2107
- LACTMED online access January 2107



To speak to a Breastfeeding Supporter call the **National Breastfeeding Helpline 0300 100 0212**

Calls to 0300 numbers cost no more than calls to UK numbers starting 01 and 02 and will be part of any inclusive minutes that apply to your provider and call package

*The Breastfeeding Network is a Company Limited by Guarantee Registered in Scotland Company No. 330639
Registered office Alexander Sloan, Chartered Accountants, 38 Cadogan Street, Glasgow, G2 7HF
The Breastfeeding Network is a Registered Scottish Charity No SC027007*

©Wendy Jones PhD, MRPharmS and the Breastfeeding Network January 2017